

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
19 April 2001 (19.04.2001)

PCT

(10) International Publication Number  
**WO 01/26468 A2**


- (51) International Patent Classification: A01N 47/00 (74) Agent: BASTIAN, Werner; Syngenta Participations AG, Intellectual Property, P.O. Box, CH-4002 Basel (CH).
- (21) International Application Number: PCT/EP00/10024 (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (22) International Filing Date: 11 October 2000 (11.10.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 09/416863 13 October 1999 (13.10.1999) CH (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (for all designated States except US): SYNGENTA PARTICIPATIONS AG [CH/CH]; CH-4058 Basel (CH).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): SENN, Robert [CH/CH]; Traubenweg 32, CH-4108 Witterswil (CH). HOFER, Dieter [CH/CH]; Oristalstrasse 46/5, CH-4410 Liestal (CH). THIEME, Thomas [DE/DE]; Birkenallee 19, 18184 Sagerheide (DE). ZANG, Larry [US/US]; 1702 Plateau Court, Highpoint, NC 27265 (US).

Published:

— Without international search report and to be republished upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHOD FOR IMPROVING PLANT GROWTH



(57) Abstract: Disclosed is a method of improving the growth of plants wherein at least one compound of formula (I), wherein A is 2-chloropyrid-5-yl, 2-methylpyrid-5-yl, 1-oxido-3-pyridinio, 2-chloro-1-oxido-5-pyridinio, 2,3-dichloro-1-oxido-5-pyridinio, tetrahydrofuran-3-yl, 5-methyl-tetrahydrofuran-3-yl or 2-chlorothiazol-5-yl group, Y is -N(R)(R<sub>2</sub>) or SR<sub>2</sub>; R is hydrogen, C<sub>1</sub>-C<sub>6</sub>alkyl, phenyl-C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl, C<sub>2</sub>-C<sub>6</sub>alkenyl or C<sub>2</sub>-C<sub>6</sub>alkynyl; R<sub>1</sub> and R<sub>2</sub> are independently of each other C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkenyl, C<sub>1</sub>-C<sub>4</sub>alkynyl, -C(=O)-CH<sub>3</sub> or benzyl; or together form a group -CH<sub>2</sub>-CH<sub>2</sub>-, -CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-, -CH<sub>2</sub>-O-CH<sub>2</sub>-, -CH<sub>2</sub>-S-CH<sub>2</sub>-, -CH<sub>2</sub>-NH-CH<sub>2</sub>- or -CH<sub>2</sub>-N(CH<sub>3</sub>)-CH<sub>2</sub>-; and X is N-NO<sub>2</sub>, N-CN or CH-NO<sub>2</sub>; or, where appropriate, a tautomer thereof, in each case in free form or in salt form, is applied to the plant or the locus thereof.

WO 01/26468 A2

## Method for Improving Plant growth

### Field of the Invention

The present invention relates to a method of improving the growth of plants comprising applying to the plants or the locus thereof at least one compound selected from the class of the neonicotinoids.

### Background of the Invention


Certain methods of improving plant growth are described in the literature. These methods are usually based on conventional fertilizing. The biological effects of those known methods are however not entirely satisfactory in the area of agriculture. There is therefore still a need to improve the growth of the plants basically for obtaining higher crop yields, as well as the reduction of the use of fertilizers needed.

### Summary of the Invention

The present invention provides a new method of improving plant growth, more specifically, a method for improving plant growth of crops such as canola (rape) seed, eggplants, rice, potatoes and soybeans, wherein at least one neonicotinoid compound is applied to the plant or the locus thereof.

### Description of Specific Embodiments

Preferred is a method of improving the growth of plants wherein at least one neonicotinoid compound of formula (I)



wherein

A is 2-chloropyrid-5-yl, 2-methylpyrid-5-yl, 1-oxido-3-pyridinio, 2-chloro-1-oxido-5-pyridinio, 2,3-dichloro-1-oxido-5-pyridinio, tetrahydrofuran-3-yl, 5-methyl-tetrahydrofuran-3-yl or 2-chlorothiazol-5-yl group,

Y is  $-N(R)(R_2)$  or  $SR_2$ ;

R is hydrogen,  $C_1$ - $C_6$ alkyl, phenyl- $C_1$ - $C_4$ alkyl,  $C_3$ - $C_6$ cycloalkyl,  $C_2$ - $C_6$ alkenyl or  $C_2$ - $C_6$ alkynyl;

$R_1$  and  $R_2$  are independently of each other  $C_1$ - $C_4$ -alkyl,  $C_1$ - $C_4$ -alkenyl,  $C_1$ - $C_4$ -alkinyl,  $-C(=O)-CH_3$  or benzyl; or together form a group  $-CH_2-CH_2-$ ,  $-CH_2-CH_2-CH_2-$ ,  $-CH_2-O-CH_2-$ ,  $-CH_2-S-CH_2-$ ,  $-CH_2-NH-CH_2-$  or  $-CH_2-N(CH_3)-CH_2-$ ; and

X is  $N-NO_2$ ,  $N-CN$  or  $CH-NO_2$ ; or, where appropriate, a tautomer thereof, in each case in free form or in salt form, is applied to the plant or the locus thereof.

The compounds (I) may be in the form of tautomers. Accordingly, hereinbefore and hereinafter, where appropriate the compound compounds (I) are to be understood to include corresponding tautomers, even if the latter are not specifically mentioned in each case.

The compounds of the formula (I) are capable of forming acid addition salts. Those salts are formed, for example, with strong inorganic acids, such as mineral acids, for example perchloric acid, sulfuric acid, nitric acid, nitrous acid, a phosphoric acid or a hydrohalic acid, with strong organic carboxylic acids, such as unsubstituted or substituted, for example halo-substituted,  $C_1$ - $C_4$ alkanecarboxylic acids, for example acetic acid, saturated or unsaturated dicarboxylic acids, for example oxalic, malonic, succinic, maleic, fumaric or phthalic acid, hydroxycarboxylic acids, for example ascorbic, lactic, malic, tartaric or citric acid, or benzoic acid, or with organic sulfonic acids, such as unsubstituted or substituted, for example halo-substituted,  $C_1$ - $C_4$ alkane- or aryl-sulfonic acids, for example methane- or p-toluene-sulfonic acid. Furthermore, compounds of formula (I) having at least one acidic group are capable of forming salts with bases. Suitable salts with bases are, for example, metal salts, such as alkali metal or alkaline earth metal salts, for example sodium, potassium or magnesium salts, or salts with ammonia or an organic amine, such as morpholine, piperidine, pyrrolidine, a mono-, di- or tri-lower alkylamine, for example ethyl-, diethyl-, triethyl- or dimethyl-propyl-amine, or a mono-, di- or tri-hydroxy-lower alkylamine, for example mono-, di- or tri-ethanolamine. In addition, corresponding internal salts may also be formed. Preference is given within the scope of the invention to agrochemically advantageous salts. In view of the close relationship between the compounds of formula (I) in free form and in the form of their salts, any reference hereinbefore or hereinafter to the free compounds of formula (I) or

to their respective salts is to be understood as including also the corresponding salts or the free compounds of formula (I), where appropriate and expedient. The same applies in the case of tautomers of compounds of formula (I) and the salts thereof. The free form is generally preferred in each case.


Preferred compounds of the formula (I) are those wherein

A is a pyrid-3-yl, 2-chloropyrid-5-yl, 2-chloro-1-oxido-5-pyridinio or 2-chlorothiazol-5-yl group; particularly a 2-chloropyrid-5-yl group or preferably a 2-chlorothiazol-5-yl group; wherein Y is  $-N(R)(R_2)$ ;

R is  $C_1$ - $C_6$ alkyl, phenyl- $C_1$ - $C_4$ alkyl,  $C_3$ - $C_4$ alkenyl or  $C_3$ - $C_4$ alkynyl; more especially  $C_1$ - $C_4$ alkyl, preferably methyl;

$R_1$  and  $R_2$  are independently of each other  $C_1$ - $C_4$ -alkyl or benzyl, or together a group  $-CH_2-CH_2-$ ,  $-CH_2-CH_2-CH_2-$ ,  $-CH_2-O-CH_2-$ ,  $-CH_2-S-CH_2-$ ,  $-CH_2-NH-CH_2-$ ,  $-CH_2-N(CH_3)-CH_2-$ , especially  $-CH_2-CH_2-$  or  $-CH_2-O-CH_2-$ , particularly  $-CH_2-O-CH_2-$ ; and X is  $N-NO_2$  or  $N-CN$ , more especially  $N-NO_2$ .

Especially preferred is a method of improving the growth of plants comprising the application to the plant or the locus thereof of an effective amount of a compound selected from the group consisting of: a compound of the formula



(Ia),

imidacloprid, clothianidin (TI-435), nitenpyram, thiacloprid, acetamiprid and MTI-446; particularly the compound of the formula (Ia) (thiamethoxam).

The compound of the formula (Ia) is known for instance from EP-A-580 553;

Imidacloprid is known from The Pesticide Manual, 11th Ed. (1997), The British Crop Protection Council, London, page 706;

Nitenpyram from The Pesticide Manual, 11th Ed. (1997), The British Crop Protection Council, London, page 880;

TI-435 from EP-A-376,279;

MTI-446 from EP-A-649'845; and  
Thiacloprid from EP-A-235'725.

Surprisingly, it has been found that the application of the compounds of the formula (I) to the plants or the locus thereof results in a quite unexpectedly enhanced plant growth. It has now been found, that the action of the compounds of the formula (I) goes far beyond their well-known pesticidal action. It has been shown, that the compounds of the formula (I) exhibit an action termed plant growth in the frame of the instant invention. Under the term plant growth there are understood various sorts of improvements of plants which are not connected to the control of pests with the said compound (I). For example such advantageous properties that may be mentioned are improved crop characteristics including: emergence, crop yields, protein content, more developed root system, tillering increase, increase in plant height, bigger leaf blade, less dead basal leaves, stronger tillers, greener leaf color, less fertilizers needed, less seeds needed, more productive tillers, earlier flowering, early grain maturity, less plant verse (lodging), increased shoot growth, improved plant vigor, and early germination; or any other advantages familiar to a person skilled in the art.

Especially preferred is the use of the said neonicotinoid compounds in a method for the improvement of the growth plants which are essentially free of insects and representatives of the order Acarina.

It has been shown, that compounds of the instant formula (I) have a good effect on the plant growth. As a rule, a good effect means at least 10% earlier emergence, crop yields, more developed root system, increase in plant height, bigger leaf blade, less fertilizers needed, less seeds needed, increased shoot growth, improved plant vigor etc.

A further aspect of the invention is a method of using a neonicotinoid compound in a method for improving the growth of plants.

A further aspect of the invention is the use of a neonicotinoid compound in a method for improving the growth of plants.

Still a further aspect of the invention is a method of using a composition comprising a neonicotinoid compound in a method for improving the growth of plants.

Crops which can be improved according to the present method include cereals, such as wheat, barley, rye, oats, rice, maize and sorghum; beet, such as sugar beet and fodder beet; fruit, for example pomes, stone fruit and soft fruit, such as apples, pears, plums, peaches, almonds, cherries and berries, e.g. strawberries, raspberries and blackberries; leguminous fruits, such as beans, lentils, peas and soybeans; oil plants, such as rape, mustard, poppy, olives, sunflowers, coconut, castor oil plants, cocoa beans and groundnuts; cucurbitaceae, such as marrows, cucumbers and melons; fibre plants, such as cotton, flax, hemp and jute; citrus fruit, such as oranges, lemons, grapefruit and mandarins; vegetables, such as spinach, lettuce, asparagus, cabbages, carrots, onions, tomatoes, potatoes and paprika; lauraceae, such as avocados, cinnamon and camphor; and also tobacco, nuts, coffee, aubergines, sugar cane, tea, pepper, vines, hops, bananas, natural rubber plants and ornamentals; especially rice, beans, soybeans, rape and potatoes.

The invention accordingly relates also to compositions comprising the compounds of the formula (I) and the use of the said compositions, such as emulsifiable concentrates, suspension concentrates, directly sprayable or dilutable solutions, coatable pastes, dilute emulsions, wettable powders, soluble powders, dispersible powders, wettable powders, dusts, granules or encapsulations in polymeric substances, which comprise at least one of the compounds according to the invention, the type of formulation being chosen in accordance with the intended objectives and the prevailing circumstances.

The compounds of the formula (I) are used in those compositions in pure form, a solid the compounds of the formula (I) being used, for example, in a specific particle size, or, preferably, together with at least one of the adjuvants customary in formulation technology, such as extenders, e.g. solvents or solid carriers, or surface-active compounds (surfactants).

Suitable formulation adjuvants are, for example, solid carriers, solvents, stabilisers, slow-release adjuvants, dyes and optionally surface-active substances (surfactants). Suitable carriers and adjuvants in this case include all substances customarily used in

crop protection products, especially in products for controlling snails and slugs. Suitable adjuvants, such as solvents, solid carriers, surface-active compounds, non-ionic surfactants, cationic surfactants, anionic surfactants and further adjuvants in the compositions used in accordance with the invention are, for example, the same as those described in EP-A-736 252; are fully incorporated by reference herein for their disclosure relating to useful formulation adjuvants.

The compositions usually contain from 0.1 to 99 %, especially from 0.1 to 95 %, of a compound of the formula (I) and from 1 to 99.9 %, especially from 5 to 99.9 %, of at least one solid or liquid adjuvant, it generally being possible for from 0 to 25 %, especially from 0.1 to 20 %, of the composition to be surfactants (in each case percentages are by weight). Whereas commercial products will preferably be formulated as concentrates, the end user will normally employ dilute formulations which have considerably lower concentrations of one or more compounds of the formula (I). Preferred formulations have especially the following composition (% = percent by weight):

Emulsifiable concentrates:

active ingredient:	1 to 90 %, preferably 5 to 20 %
surface-active agent:	1 to 30 %, preferably 10 to 20 %
liquid carrier:	5 to 94 %, preferably 70 to 85 %

Dusts:

active ingredient:	0.1 to 10 %, preferably 0.1 to 1 %
solid carrier:	99.9 to 90 %, preferably 99.9 to 99 %

Suspension concentrates:

active ingredient:	5 to 75 %, preferably 10 to 50 %
water:	94 to 24 %, preferably 88 to 30 %
surface-active agent:	1 to 40 %, preferably 2 to 30 %

Wettable powders:

active ingredient:	0.5 to 90 %, preferably 1 to 80 %
surface-active agent:	0.5 to 20 %, preferably 1 to 15 %
solid carrier:	5 to 95 %, preferably 15 to 90 %

Granules:

active ingredient:	0.5 to 30 %, preferably 3 to 15 %
solid carrier:	99.5 to 70 %, preferably 97 to 85 %

Injection solution:

active ingredient:	0.1 to 10 %, preferably 0.5 to 5 %
non-ionic surfactant:	0.1 to 30 %, preferably 0.5 to 10 %
mixture of ethanol and propylene glycol:	60 to 99 %, preferably 85 to 90 %

Injection suspension (aqueous or oily):

active ingredient:	0.1 to 20 %, preferably 1 to 10 %
non-ionic surfactant:	0.1 to 20 %, preferably 1 to 10 %
water or vegetable oil:	60 to 99 %, preferably 85 to 95 %

The compositions according to the invention are prepared in known manner: in the absence of adjuvants, for example, by grinding, sieving and/or compressing a solid compound of the formula (I), for example to a specific particle size, and, in the presence of at least one adjuvant, for example, by intimately mixing and/or grinding the compound of the formula (I) with the adjuvant(s). The invention relates also to those methods of preparing the compositions according to the invention and to the use of compounds I in the preparation of such compositions.

The invention relates also to the methods of applying the compositions of the type mentioned, such as spraying, atomising, dusting, coating, dressing, scattering or pouring, which are chosen in accordance with the intended objectives and the prevailing circumstances, and to the use of the compositions for the improvement of the plants of the type mentioned. Typical rates of concentration are from 0.1 to 1000 ppm, preferably from 0.1 to 500 ppm, of compound of the formula (I). The rates of application per hectare are generally from 1 to 2000 g of compound of the formula (I) per hectare, especially from 1 to 1000 g/ha, preferably from 5 to 600 g/ha.

A preferred method of application is application to the leaves of the plants (foliar application), the frequency and rate of application depending on the desired



improvement of the crop plant in question. The compound of the formula (I) may, however, also penetrate the plants through the root system (systemic action) as a result of impregnation of the locus of the plant with a liquid formulation or by incorporation of the compound of the formula (I) in solid form, for example in the form of granules, in the locus of the plant, for example in the soil (soil application). In the case of paddy rice crops, such granules may be applied in metered amounts to the flooded rice field.

In one embodiment, commercial products will preferably be formulated as concentrates whereas the end user will normally use dilute formulations.

The compositions according to the invention are also suitable for the treatment of plant propagation material, including genetically modified propagation material, e.g. seed, such as fruit, tubers or grains, or plant cuttings. The propagation material may be treated with the composition before planting, for example seed may be dressed before sowing. The compounds according to the invention may also be applied to seed grains (coating) either by impregnating the grains with a liquid formulation or by coating them with a solid formulation. The composition may also be applied to the planting site when the propagation material is being planted, for example may be applied to the seed furrow during sowing. The invention relates also to that method of treating plant propagation material and to the plant propagation material so treated.

The compounds of formula (I) are normally applied to plant propagation material in the form of compositions, but also can be applied to the seed or to the locus of propagation thereof (such as a furrow), simultaneously or in succession, with further compounds. These further compounds can be fertilizers or micronutrient donors or other preparations that influence plant growth. They can also be selective pesticides or mixtures of several of these preparations, if desired together with further carriers, surfactants or application-promoting adjuvants customarily employed in the art of formulation.

In connection with the treatment of plant propagation material, favorable rates of application are in general 0.0005 to not more than 1 kg, in particular 0.01–0.8 kg, more particularly 0.1–0.5 kg of one or more compounds of the formula (I) per 100 kg of material to be protected. However, the application conditions depend essentially on the

nature (surface area, consistency, moisture content) of the material and on its environmental factors. Accordingly, within these ranges, those skilled in the art will choose, on the basis of their general body of knowledge and, where appropriate, a few experiments, doses which are non-phytotoxic but effective for improving the plant growth.

The techniques of seed treatment application are well known to those skilled in the art, and they may be used readily in the context of the present invention. The compounds of the formula (I) can be formulated and applied as a slurry, a solid seed coating, a soak, or as a dust on the surface of the seed. There also may be mentioned, e.g., film-coating or encapsulation. The coating processes are well known in the art, and employ, for seeds, the techniques of film-coating or encapsulation, or for the other multiplication products, the techniques of immersion. Needless to say, the method of application of the compounds to the seed may be varied and the invention is intended to include any technique which is to be used.

A preferred method of applying the mixture to the plant propagation material according to the invention consists in spraying or wetting the plant propagation material with a liquid preparation, or mixing the plant material with a solid preparation of the compounds of the formula (I).

The compounds of this invention may be formulated or mixed in the seed treater tank or combined on the seed by overcoating with other seed treating agents. The agents to be mixed with the compounds of this invention may be for the control of pests, or further modification of growth, nutrition, or for the control of plant diseases.

Formulation Examples (% = per cent by weight)

The examples which follow are intended to illustrate and not limit the invention, "compound of the formula (I)" being understood as meaning one or several of the compounds of the formula (I).

Example F1: Emulsifiable concentrates	a)	b)	c)
Compound of the formula (I)	25%	40%	50%
calcium dodecylbenzenesulfonate	5%	8%	6%
castor oil polyethylene glycol ether	5%	-	-
Tributylphenol polyethylene glycol ether	-	12%	4%
Cyclohexanone	-	15%	20%
xylene mixture	65%	25%	20%

Emulsions of any desired concentration can be prepared from this concentrate by dilution with water, and can be employed in crop protection and in seed treatment applications.

Example F2: Dusts	a)	b)
Compound of the formula (I)	5%	8%
Talc	95%	-
Kaolin	-	92%

Ready-to-use dusts are obtained by mixing the compounds of the formula (I) with the carrier and grinding the mixture in a suitable mill. Such powders can be used for dry-dressing seeds.

Example F3: Wettable powders	a)	b)	c)
Compound of the formula (I)	25%	50%	75%
sodium lignosulfonate	5%	5%	-
sodium laurylsulfate	3%	-	5%
sodium diisobutylphenathalenesulfonate	-	6%	10%
octylphenol polyethylene glycol ether	-	2%	-
highly dispersed silicic acid	5%	10%	10%
kaolin	62%	27%	-

The compound of the formula (I) is mixed thoroughly with the additives and the mixture is ground thoroughly in a suitable mill affording wettable powders which can be diluted

with water to give suspensions of any desired concentration. Such slurries can be used for carrying out furrow treatments on prior to planting crops of plants and also for wet- or moist-dressing material which can be propagated, for example oil seeds or tubers of plants.

Example F4: Suspoemulsions	a)
Compound of the formula (I)	22.5%
sulfated nonylphenol (polyoxyethylene condensate)	0.1%
phosphated tristerylphenol (polyoxyethylene condensate)	4%
sodium lignosulfonate (polyoxyethylene condensate)	2%
NaOH (50%)	0.1%
silicone defoaming agent	0.1%
pigment(s)	9.5%
Glycerin	20%
xanthan gum	0.2%
Water	41.5%

This formulation is suitable for mixtures of solid and liquid compounds of the formula (I). The solid compounds of the formula (I) are mixed thoroughly with a portion of the emulsifiers and water and the mixture is ground thoroughly in a suitable mill. Another portion of the emulsifiers and water are mixed with the liquid compounds of the formula (I). The two mixtures are combined along with any other inert ingredients (such as pigments, thickeners, etc.) that are to be used in the formulation. Such suspoemulsions can be used for carrying out in furrow treatments prior to planting crops of plants and also for wet- or moist-dressing material which can be propagated, for example oil seeds or tubers of plants.

Biological Examples (% = per cent by weight unless otherwise indicated)

The examples which follow are intended to illustrate and not limit the invention.

Example B1: Emergence

Canola seed is treated with the composition of the invention containing Thiamethoxam, at a rate of 400 g Thiamethoxam per 100 kg seed, and is seeded following procedures which correspond to conditions found in practice. Untreated seeds from the same origin are used for comparison purposes. Emergence (plants per meter) is evaluated. In the case of treated seeds, about 20% more plants per meter emerge than in the case of untreated plants.

Example B2 / B3: Vigor of Rice (*Oryza sativa* L. cv. Nihonbare) and Eggplant (*Solanum melongena* L. cv. Marfa)

Plants are grown in a commercially available pathogen-free soil mixture. 2 weeks (eggplants 3 weeks) after planting seedlings are transplanted into 600 ml pots and drench treated with 10 mg Thiamethoxam in 20 ml water per pot. They were fertilised once or twice a week. Up to 10 weeks after treatment, fresh- and dry-weight of shoots and roots and numbers of leaves are determined. Protein content is measured by the method of Bradford (1976, Anal. Biochem., 72, 248-254) with BSA (bovine serum albumin) as standard: Homogenisation of leaf material in liquid N<sub>2</sub> in a mortar. Extraction in phosphate buffer (pH 7.5, 0.1M, 2/1 vol/weight). After centrifugation supernatant is separated from pellet and stored at -20°C until protein determination. For comparison of treated and check, mg protein/g fresh weight is calculated.

Tables B2 and B3: Differences between treated plants and Check given as factors (e.g. Fresh weight of treated/ Fresh weight of check). Factors >1 indicate an enhancement of vigor of treated plants. FW: fresh weight; DW: dry weight; Protein ((mg/g treated plant)/(mg/g check))

Table B2: Rice

Week	Size	Shoot		Root		Protein
		FW	DW	FW	DW	
1	1.0	1.0	1.0	1.0	0.8	1.1
2	1.0	1.0	1.1	0.8	1.1	1.0
3	1.0	1.2	1.3	0.6	1.3	1.0

13

4	1.1	1.3	1.2	1.5	1.4	1.0
5	1.1	1.2	1.1	1.0	0.9	1.0
6	1.2	1.3	1.1	1.6	1.5	1.0
7	1.1	1.3	1.2	1.3	1.3	1.1
8	1.1	1.4	1.4	1.8	1.5	1.0
9	1.2	1.2	1.2	1.0	1.0	0.9
10	1.3	1.3	1.3	2.0	2.0	0.7

Table B3: Eggplant

Week	Size	Shoot		Root		Leaf	Protein
		FW	DW	FW	DW	No	
1	1.0	0.9	0.9	1.0	1.0		1.0
2	1.1	0.7	1.1	1.2	1.2		1.1
3	1.1	1.2	1.1	1.1	1.1		1.0
4	1.0	1.1	1.1	1.0	1.0		1.6
5	1.0	1.0	1.1	0.9	1.0	1.1	1.0
6	1.3	1.3	1.3	0.9	0.9	1.3	1.1
7	1.2	1.1	1.1	0.8	0.7	1.3	1.4
8	1.2	0.7	0.9	0.6	0.6	1.2	1.4
9	1.4	0.7	1.1	0.9	1.1	1.6	2.5
10	1.4	1.1	1.1	1.1	1.3	1.4	1.3

Since the soil used does not contain pathogens and the biological activity can be assumed as low, the effects are caused by direct growth stimulation and not by side effects of Thiamethoxam against soil organisms.

Example B4: Yield


Potato tubers are treated with the composition of the invention containing Imidacloprid, at a rate of 500 g Imidacloprid per 100 kg seed and are planted following procedures which correspond to conditions found in practice. The crop is harvested from the field at maturity. Untreated tubers from the same origin are used for comparison purposes. Crop yield is evaluated and found to be significantly higher in the case of treated potato tubers than with untreated tubers.

In summary, it is seen that this invention provides a new method for improving the plant growth. Variations may be made in proportions, procedures and materials without departing from the scope of the invention as defined by the following claims.

**We claim:**

1. A method of improving the growth of plants comprising applying to the plants or the locus thereof at least one compound selected from the class of the neonicotinoids.

2. A method according to claim 1,  
wherein at least one compound of formula (I)



wherein

A is 2-chloropyrid-5-yl, 2-methylpyrid-5-yl, 1-oxido-3-pyridinio, 2-chloro-1-oxido-5-pyridinio, 2,3-dichloro-1-oxido-5-pyridinio, tetrahydrofuran-3-yl, 5-methyl-tetrahydrofuran-3-yl or 2-chlorothiazol-5-yl group,

Y is  $-\text{N}(\text{R})(\text{R}_2)$  or  $\text{SR}_2$ ;

R is hydrogen,  $\text{C}_1$ - $\text{C}_6$ alkyl, phenyl- $\text{C}_1$ - $\text{C}_4$ alkyl,  $\text{C}_3$ - $\text{C}_6$ cycloalkyl,  $\text{C}_2$ - $\text{C}_6$ alkenyl or  $\text{C}_2$ - $\text{C}_6$ alkynyl;

$\text{R}_1$  and  $\text{R}_2$  are independently of each other  $\text{C}_1$ - $\text{C}_4$ -alkyl,  $\text{C}_1$ - $\text{C}_4$ -alkenyl,  $\text{C}_1$ - $\text{C}_4$ -alkinyl,  $-\text{C}(=\text{O})-\text{CH}_3$  or benzyl; or together form a group  $-\text{CH}_2-\text{CH}_2-$ ,  $-\text{CH}_2-\text{CH}_2-\text{CH}_2-$ ,  $-\text{CH}_2-\text{O}-\text{CH}_2-$ ,  $-\text{CH}_2-\text{S}-\text{CH}_2-$ ,  $-\text{CH}_2-\text{NH}-\text{CH}_2-$  or  $-\text{CH}_2-\text{N}(\text{CH}_3)-\text{CH}_2-$ ; and

X is  $\text{N}-\text{NO}_2$ ,  $\text{N}-\text{CN}$  or  $\text{CH}-\text{NO}_2$ ; or, where appropriate, a tautomer thereof, in each case in free form or in salt form, is applied to the plant or the locus thereof.

3. A method according to claim 2, wherein Thiamethoxam is applied to the plants or the locus thereof.

4. A method according to claim 2, wherein Imidacloprid is applied to the plants or the locus thereof.

5. A method of using a compound of the formula (I) for improving the growth of plants, wherein the said compound of the formula (I) is applied to the plants or the locus thereof.



6. The use of a compound of a neonicotinoid compound in a method as claimed in claim 1.
7. The use of a composition comprising at least one neonicotinoid compound in a method as claimed in claim 1.
8. A method of using a composition comprising a neonicotinoid compound in a method as claimed in claim 1 for improving the growth of plants.

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
19 April 2001 (19.04.2001)


PCT

(10) International Publication Number  
**WO 01/26468 A3**

- (51) International Patent Classification<sup>7</sup>: A01N 51/00, 47/44, 43/78, 43/40, 43/08, A01C 1/06
- (21) International Application Number: PCT/EP00/10024
- (22) International Filing Date: 11 October 2000 (11.10.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
09/416863 13 October 1999 (13.10.1999) CH
- (71) Applicant (for all designated States except US): SYNGENTA PARTICIPATIONS AG [CH/CH]; CH-4058 Basel (CH).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): SENN, Robert [CH/CH]; Traubenweg 32, CH-4108 Witterswil (CH). HOFER, Dieter [CH/CH]; Oristalstrasse 46/5, CH-4410 Liestal (CH). THIEME, Thomas [DE/DE]; Birkenallee 19, 18184 Sagerheide (DE). ZANG, Larry [US/US]; 1702 Plateau Court, Highpoint, NC 27265 (US).
- (74) Agent: BASTIAN, Werner; Syngenta Participations AG, Intellectual Property, P.O. Box, CH-4002 Basel (CH).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— with international search report
- (88) Date of publication of the international search report:  
1 November 2001
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/26468 A3

(54) Title: METHOD FOR IMPROVING PLANT GROWTH



(57) Abstract: Disclosed is a method of improving the growth of plants wherein at least one compound of formula (I), wherein A is 2-chloropyrid-5-yl, 2-methylpyrid-5-yl, 1-oxido-3-pyridinio, 2-chloro-1-oxido-5-pyridinio, 2,3-dichloro-1-oxido-5-pyridinio, tetrahydrofuran-3-yl, 5-methyl-tetrahydrofuran-3-yl or 2-chlorothiazol-5-yl group, Y is -N(R)(R<sub>2</sub>) or SR<sub>2</sub>; R is hydrogen, C<sub>1</sub>-C<sub>6</sub>alkyl, phenyl-C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl, C<sub>2</sub>-C<sub>6</sub>alkenyl or C<sub>2</sub>-C<sub>6</sub>alkynyl; R<sub>1</sub> and R<sub>2</sub> are independently of each other C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkenyl, C<sub>1</sub>-C<sub>4</sub>alkynyl, -C(=O)-CH<sub>3</sub> or benzyl; or together form a group -CH<sub>2</sub>-CH<sub>2</sub>-, -CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-, -CH<sub>2</sub>-O-CH<sub>2</sub>-, -CH<sub>2</sub>-S-CH<sub>2</sub>-, -CH<sub>2</sub>-NH-CH<sub>2</sub>- or -CH<sub>2</sub>-N(CH<sub>3</sub>)-CH<sub>2</sub>-; and X is N-NO<sub>2</sub>, N-CN or CH-NO<sub>2</sub>; or, where appropriate, a tautomer thereof, in each case in free form or in salt form, is applied to the plant or the locus thereof.

# INTERNATIONAL SEARCH REPORT

Internat I Application No  
PCT/EP 00/10024

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A01N51/00 A01N47/44 A01N43/78 A01N43/40 A01N43/08  
A01C1/06

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A01N A01C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>PALUMBO J C ET AL: "IMIDACLOPRID DOES NOT ENHANCE GROWTH AND YIELD OF MUSKMELON IN THE ABSENCE OF WHITEFLY"</p> <p>HORTSCIENCE, AMERICAN SOCIETY OF HORTICULTURAL SCIENCE, ALEXANDRIA, VA, US, vol. 30, no. 5, August 1995 (1995-08), pages 997-999, XP000985863</p> <p>ISSN: 0018-5345</p> <p>page 997, column 2, line 8-11</p> <p>page 998, column 2, paragraph 2 -column 3, paragraph 2</p> <p>page 999, column 1, paragraph 2</p> <p>---</p> <p>-/--</p>	1-8

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*G\* document member of the same patent family

Date of the actual completion of the international search

9 May 2001

Date of mailing of the international search report

31/05/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Klaver, J

# INTERNATIONAL SEARCH REPORT

Internal Application No  
PCT/EP 00/10024

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>S. J. FLEISCHER, M. D. ORZOLEK, D. DE MACKIEWICZ &amp; L. OTJEN: "Imidacloprid effects on <i>Acalymna vittata</i> (Coleoptera: Chrysomelidae) and bacterial wilt in cantaloupe." J. ECON. ENTOMOL., vol. 91, no. 4, 1998, pages 940-949, XP002166302 Entomol. Soc. of America page 943, column 1, paragraph 2 -page 944, column 1, paragraph 2 page 948, column 1, paragraph 1 ---</p>	1-8
X	<p>J. C. PALUMBO, D. L. KERNS, C. E. ENGLE, C. A. SANCHEZ &amp; M. WILCOX: "Imidacloprid formulation and soil placement effects on colonization by seetpotato whitefly (Homoptera: Aleyrodidae): headsize and incidence of chlorosis in lettuce." J. ECON. ENTOMOL., vol. 89, no. 3, 1996, XP002166303 Entomol. Soc. of America page 738, column 1, paragraph 3 -column 2, paragraph 1 page 741, column 1, paragraph 2 ---</p>	1-8
X	<p>K. S. PIKE, G. L. REED, G. T. GRAF &amp; D. ALLISON: "Imidacloprid systemic seed treatment effects on germination, plant growth, yiled and compatibility with seed-applied fungicides and control of Russian Wheat Aphid in wheat and barley." GREAT PLAINS AGRIC.COUNCIL PUBL.: PROC. OF 5TH RUSSIAN WHEAT APHID CONFERENCE, JAN. 26-28, 1992, no. 142, 1992, XP000995492 Fort Worth, Texas the whole document ---</p>	1-8
Y	<p>US 5 876 739 A (CHEN YUGUANG ET AL) 2 March 1999 (1999-03-02) column 4, line 25-32 ---</p>	1-8
Y	<p>EP 0 113 070 A (AMERICAN CYANAMID CO) 11 July 1984 (1984-07-11) page 1, line 10 -page 3, line 20 page 7, line 3-29 page 8, line 12-20 -----</p>	1-8

# INTERNATIONAL SEARCH REPORT

Information on patent family members

Interna. Application No

PCT/EP 00/10024

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5876739 A	02-03-1999	US 5849320 A	15-12-1998
EP 0113070 A	11-07-1984	AT 31920 T	15-01-1988
		AU 565227 B	10-09-1987
		AU 2251583 A	28-06-1984
		AU 7556787 A	22-10-1987
		BR 8306966 A	24-07-1984
		DD 225988 A	14-08-1985
		DE 3345281 A	20-06-1984
		DK 585083 A	21-06-1984
		EP 0189579 A	06-08-1986
		ES 528171 D	16-11-1985
		ES 8601115 A	16-02-1986
		ES 543756 D	16-10-1986
		ES 8700229 A	01-01-1987
		ES 543758 D	16-01-1986
		ES 8604120 A	01-06-1986
		FI 834676 A	03-06-1984
		FI 882004 A	28-04-1988
		HU 195637 B	28-06-1988
		JP 59118757 A	09-07-1984
		NZ 206556 A	12-02-1988
		PH 21256 A	31-08-1987
		US 4944788 A	31-07-1990
		US 4594092 A	10-06-1986
		US 4804780 A	14-02-1989
		ZA 8309399 A	29-08-1984

## CORRECTED VERSION

(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
19 April 2001 (19.04.2001)


PCT

(10) International Publication Number  
**WO 2001/026468 A3**

- (51) International Patent Classification<sup>7</sup>: **A01N 51/00**, 47/44, 43/78, 43/40, 43/08, A01C 1/06
- (21) International Application Number: PCT/EP2000/010024
- (22) International Filing Date: 11 October 2000 (11.10.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
09/416,863 13 October 1999 (13.10.1999) US
- (71) Applicant (for all designated States except US): SYNGENTA PARTICIPATIONS AG [CH/CH]; CH-4058 Basel (CH).
- (72) Inventors; and  
(75) Inventors/Applicants (for US only): SENN, Robert [CH/CH]; Traubenweg 32, CH-4108 Witterswil (CH). HOFER, Dieter [CH/CH]; Oristalstrasse 46/5, CH-4410 Liestal (CH). THIEME, Thomas [DE/DE]; Birkenallee 19, 18184 Sagerheide (DE). ZANG, Larry [US/US]; 1702 Plateau Court, Highpoint, NC 27265 (US).
- (74) Agent: BASTIAN, Werner; Syngenta Participations AG, Intellectual Property, P.O. Box, CH-4002 Basel (CH).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— with international search report
- (88) Date of publication of the international search report: 1 November 2001
- (48) Date of publication of this corrected version: 19 February 2004
- (15) Information about Correction:  
see PCT Gazette No. 08/2004 of 19 February 2004, Section II
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 2001/026468 A3

(54) Title: METHOD FOR IMPROVING PLANT GROWTH



(57) Abstract: Disclosed is a method of improving the growth of plants wherein at least one compound of formula (I), wherein A is 2-chloropyrid-5-yl, 2-methylpyrid-5-yl, 1-oxido-3-pyridinio, 2-chloro-1-oxido-5-pyridinio, 2,3-dichloro-1-oxido-5-pyridinio, tetrahydrofuran-3-yl, 5-methyl-tetrahydrofuran-3-yl or 2-chlorothiazol-5-yl group, Y is -N(R)(R<sub>2</sub>) or SR<sub>2</sub>; R is hydrogen, C<sub>1</sub>-C<sub>6</sub>alkyl, phenyl-C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>3</sub>-C<sub>6</sub>cycloalkyl, C<sub>2</sub>-C<sub>6</sub>alkenyl or C<sub>2</sub>-C<sub>6</sub>alkynyl; R<sub>1</sub> and R<sub>2</sub> are independently of each other C<sub>1</sub>-C<sub>4</sub>alkyl, C<sub>1</sub>-C<sub>4</sub>alkenyl, C<sub>1</sub>-C<sub>4</sub>alkinyl, -C(=O)-CH<sub>3</sub> or benzyl; or together form a group -CH<sub>2</sub>-CH<sub>2</sub>-, -CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-, -CH<sub>2</sub>-O-CH<sub>2</sub>-, -CH<sub>2</sub>-S-CH<sub>2</sub>-, -CH<sub>2</sub>-NH-CH<sub>2</sub>- or -CH<sub>2</sub>-N(CH<sub>3</sub>)-CH<sub>2</sub>-; and X is N-NO<sub>2</sub>, N-CN or CH-NO<sub>2</sub>; or, where appropriate, a tautomer thereof, in each case in free form or in salt form, is applied to the plant or the locus thereof.